



COVID-19 Consent & Medical Eligibility

For Comirnaty® or Comirnaty® RTU, Comirnaty® BA.1 or Comirnaty® BA.4-5 COVID-19 vaccines and Spikevax® or Spikevax® BA.1 COVID-19 vaccines



Complete this part for the person being vaccinated (PLEASE USE BLOCK CAPITALS)

Name: [] Date of Birth: [] [] []

Please answer the following questions with a yes or no answer

1. Has this person ever had a serious allergic reaction (anaphylaxis) that needed medical treatment:

I) after having a previous dose of the Spikevax® or Spikevax® BA.1 or Comirnaty® or Comirnaty® RTU or Comirnaty® BA.1 or Comirnaty® BA.4-5 COVID-19 vaccine, OR

Yes No [] []

II) to any of the vaccine ingredients, including polyethylene glycol known as PEG?

If yes, they cannot get this vaccine. If no, GO TO NEXT QUESTION.

1a. Have they ever had a serious allergic reaction (anaphylaxis) to Trometamol (an ingredient in contrast dye used in MRI radiological studies)?

Yes No [] []

If yes, they cannot get the Spikevax® or Spikevax® BA.1 or Comirnaty® RTU or Comirnaty® BA.1 or Comirnaty® BA.4-5 COVID-19 vaccine. But they can have a different vaccine. If no, GO TO NEXT QUESTION.

2. Have they ever had a serious allergic reaction (anaphylaxis):

I) after taking multiple different medications, with no reason known for the reaction. This may mean they are allergic to polyethylene glycol (PEG) OR

Yes No [] []

II) after having a vaccine or a medicine that contains polyethylene glycol (PEG), OR

III) for unexplained reasons. This may mean they are allergic to polyethylene glycol (PEG)?

If yes, they cannot get this vaccine, they may need specialist advice. Talk to the vaccination team. If no, GO TO NEXT QUESTION.

3. Have they ever had Mastocytosis (rare condition caused by an excess number of mast cells gathering in the body's tissues)

Yes No [] []

If yes, they can still get the vaccine, BUT, they should be observed for 30 minutes after they are vaccinated. If no, GO TO NEXT QUESTION.

4. Have they had myocarditis (inflammation of the heart muscle) after having a previous dose of the Spikevax® or Spikevax® BA.1 or Comirnaty® or Comirnaty® RTU or Comirnaty® BA.1 or Comirnaty® BA.4-5 COVID-19 vaccine?

Yes No [] []

If yes, they cannot get this vaccine. If no, GO TO NEXT QUESTION.

5. Have they had pericarditis (inflammation of the lining around the heart) after having a previous dose of the Spikevax® or Spikevax® BA.1 or Comirnaty® or Comirnaty® RTU or Comirnaty® BA.1 or Comirnaty® BA.4-5 COVID-19 vaccine?

Yes No [] []

If yes, go to Question 5a. If no, this person can be vaccinated today. GO TO NEXT QUESTION.

5a. Since they had pericarditis (inflammation of the lining around the heart) after a previous dose of the Spikevax® or Spikevax® BA.1 or Comirnaty® or Comirnaty® RTU or Comirnaty® BA.1 or Comirnaty® BA.4-5 COVID-19 vaccine, a specialist doctor must approve they get this vaccine.

Yes No [] []

Has their COVID-19 vaccination been approved by a specialist doctor?

If yes, GO TO NEXT QUESTION. If no, they cannot get this vaccine.

Question 6 for children aged 5-11 years only

6. Has this child had multisystem inflammatory syndrome also called MIS-C (a rare syndrome usually treated in hospital) after a COVID-19 infection?

Yes No [] []

If yes, please answer question 6a and 6b

6a. Has this child clinically recovered from MIS-C?

Yes No [] []

If yes, this child can be vaccinated today. If no, this child cannot be vaccinated today.

6b. Has it been over 90 days since this child was diagnosed with MIS-C?

Yes No [] []

If yes, this child can be vaccinated today. If no, this child cannot be vaccinated today

7. Have they had the monkeypox or smallpox vaccine (Imvanex or Jynneos) in the last 4 weeks?

Yes No [] []

If yes, they cannot get this vaccine today. They need to wait 4 weeks after getting these vaccines before getting a COVID-19 vaccine.



COVID-19 Consent & Medical Eligibility

For Comirnaty® or Comirnaty® RTU, Comirnaty® BA.1 or Comirnaty® BA.4-5 COVID-19 vaccines and Spikevax® or Spikevax® BA.1 COVID-19 vaccines



Please answer the following questions with a yes or no answer

8. If receiving your first or second dose of a COVID-19 vaccine, have you been diagnosed with COVID-19 (with a PCR or Antigen test) within the last four weeks?

If yes and receiving a first or second dose of a COVID-19 vaccine, you should delay getting a vaccine until you have recovered from COVID-19 and it has been at least four weeks since you tested positive or developed symptoms, or four weeks from your first positive PCR if you did not have symptoms. If no, GO TO NEXT QUESTION.

Yes No

9. Has this person tested positive (with a PCR or antigen test) for COVID-19 in the last 4 months?

If yes, they should delay getting an additional or booster dose of COVID-19 for 4 months after diagnosis or onset of symptoms. If no, GO TO NEXT QUESTION.

Yes No

10. Does this person have a bleeding disorder or are they on anticoagulation therapy?

If yes, they can still get a vaccine if they have a bleeding disorder or take anticoagulation medicines. But tell their vaccinator about their condition.

Yes No

11. Is this person 29 years of age or younger?

If yes, they can be offered a dose of Comirnaty® or Comirnaty® RTU or Comirnaty® BA.1 or Comirnaty® BA.4-5. If no, they can be offered either a dose of Comirnaty® or Comirnaty® RTU or Comirnaty® BA.1 or Comirnaty® BA.4-5 COVID-19 vaccines and Moderna (Spikevax®) or Spikevax® BA.1

Yes No

12. For Moderna (Spikevax®) only. Has this person ever been diagnosed with capillary leak syndrome?

If yes, GO TO QUESTION 12a.

Yes No

12a. Has this person had a discussion on the risks and benefits of this vaccine with their GP or specialist doctor and they have approved the vaccine for them?

Yes No

13. Is this person pregnant and attending for a second booster or subsequent booster?

If yes, GO TO QUESTION 13a.

Yes No

13a. Are you either under 16 weeks or already had a first booster in this pregnancy?

If yes, you are ineligible. You need to wait until you are 16 weeks pregnant before you can get the second or subsequent booster unless you are otherwise eligible with other risk factors (see latest NIAC guidelines).

Yes No

You only need one booster dose per pregnancy.

For people aged 16 years and older

One of these options is appropriate when establishing consent (please tick as appropriate)

- 1. The individual has consented to vaccination for COVID-19 and has been provided with written information, **OR**
- 2. The individual does not agree with COVID-19 vaccination and should not be vaccinated, **OR**
- 3. The individual cannot consent and they are being vaccinated for COVID-19 according to their benefit and will and preference, **AND**

The above is recorded in their healthcare record and includes information about any consultation that has taken place to help determine their will and preference.

For people aged 15 years and younger

Please note only a parent or legal guardian can consent or refuse consent for people aged 15 years and younger.

I confirm that I am authorised to give consent on behalf of the above named young person.

I understand I am giving consent for the administration of a dose or for the primary course consent for the administration of two or three doses of COVID-19 at the appropriate interval.

Signature Date (DD/MM/YYYY)

Name (Please print) (Please tick) Parent Legal Guardian Self

This Young person assents to receiving the vaccine (Please tick)

FOR OFFICE USE ONLY

Name of Vaccinator Registration Number / PIN / MCRN